arXiv:1405.6072v2 [math.GT] 8 Dec 2015

VIRTUAL COVERS OF LINKS

MICAH CHRISMAN

Abstract. We use virtual knot theory to detect the non-invertibility of some classical links in $\mathbb{S}^3$. These links appear in the study of virtual covers. Briefly, a virtual cover associates a virtual knot v to a knot K in a 3-manifold N, under certain hypotheses on K and N. Virtual covers of links in $\mathbb{S}^3$ come from taking K to be in the complement N of a fibered link J. If $J \sqcup K$ is invertible and K is "close to" a fiber of J, then v satisfies a symmetry condition to which some virtual knot polynomials are sensitive. We also consider an application of virtual covers to finite-type invariants. Additionally, we discuss virtual covers of links $J \sqcup K$ where J is not fibered but is virtually fibered (in the sense of W. Thurston).

It is well-known that neither the HOMFLYPT polynomial nor any quantum invariants can distinguish a link in $\mathbb{S}^3$ from its inverse. Other methods can do it, but they are difficult [24, 10]. The situation for virtual knots is better: easily computable asymmetry sensitive virtual knot invariants exist. Recently [4, 17], the technique of virtual covers was developed. It naturally associates virtual knots to *some* two component links. These observations will be exploited to give simple proofs of non-invertibility for *some* links in $\mathbb{S}^3$ (see Section 1.6).

To do this, we build on [4] and establish a condition on links in $\mathbb{S}^3$ so that the associated virtual knot functions essentially as a link invariant (see Section 1.5). This result has other uses. We show (Section 2.2) that if the Vassiliev invariants separate ordered oriented links in $\mathbb{S}^3$, then they also separate knots in thickened c.c.o surfaces with non-empty boundary, up to equivalence of virtual knots. Further applications will be considered in [3].

These diverse applications suggest that it is desirable to extend virtual covers from *some* links to *some more* links. In [4], two component links of the form $J \sqcup K$ with J fibered and $\mathrm{lk}(J, K) = 0$ were considered. This generalizes easily to $m + 1$ component links $J \sqcup K$ with J a fibered m component link and K "close to" a fiber of J (see Section 1.4). Section 3 considers the case that J is not necessarily fibered but that $\mathbb{S}^3 \backslash J$ has a finite-index cover by $\mathbb{S}^3 \backslash \tilde{J}$, where $\tilde{J}$ is a fibered link. Such links J are examples of *virtually fibered links*, a class of links that conjecturally contains all hyperbolic links [22].

1. Virtual Covers and Non-Invertibility of Links

1.1. **Review: Virtual Knots.** The four models of virtual knots are:

(1) virtual knot diagrams modulo the extended Reidemeister moves [13],
(2) Gauss diagrams modulo diagrammatic Reidemeister moves [8],
(3) abstract knot diagrams modulo Kamada–Kamada equivalence [2, 11], and
(4) knots in thickened surfaces modulo stabilization and destabilization [18].

2000 *Mathematics Subject Classification.* 57M25, 57M27.
Key words and phrases. virtual knots, non-invertible links, Vassiliev filtration, virtually fibered links.

There is a one-to-one correspondence between the equivalence classes of each interpretation. If we have a representative R of one of the models (2), (3), or (4), we will denote by $\kappa(R)$ the corresponding equivalence class of virtual knot diagrams from model (1). The four models for a single virtual knot are shown below.



We use "$\leftrightharpoons$" to denote equivalence of virtual knots. If a virtual knot has a diagram whose crossings are all classical, then it is said to be a *classical knot*. If two classical knots are equivalent as virtual knots, then they are equivalent as knots in $\mathbb{S}^3$ [8]. A virtual knot is classical if and only if it can be represented as a diagram on a compact planar surface.

1.2. **Virtual Covers.** Let M be a smooth connected oriented 3-manifold and K^M a smooth oriented knot in $\text{int}(M)$. Knots in M are considered equivalent up to ambient isotopy in M. If $\partial M \neq \emptyset$, we assume that the isotopy acts as the identity on ∂M. We write $K_1^M \leftrightharpoons K_2^M$ if K_1^M and K_2^M are equivalent in M.

Let N be a compact connected oriented (c.c.o) 3-manifold having a regular orientation preserving covering space $\Pi : \Sigma \times \mathbb{R} \to N$, where Σ is a c.c.o smooth 2-manifold. If K^N is a knot in N and $\mathfrak{k}^{\Sigma\times\mathbb{R}}$ is a knot in $\Sigma \times \mathbb{R}$ satisfying $\Pi(\mathfrak{k}) = K$, we call $\mathfrak{k}^{\Sigma\times\mathbb{R}}$ a *lift by* Π of K^N. A *virtual cover* of K^N is choice $\mathfrak{k}^{\Sigma\times\mathbb{R}}$ of lift by Π for K^N, denoted $(\mathfrak{k}^{\Sigma\times\mathbb{R}}, \Pi, K^N)$. The oriented virtual knot $\upsilon = \kappa(\mathfrak{k}^{\Sigma\times\mathbb{R}})$ is called the *associated virtual knot*.



Viewing K as a closed path at x_0 in N, regularity implies there is a one-to-one correspondence between lifts by Π and elements of $\Pi^{-1}(x_0)$. If K^N has a lift by Π, let $\Upsilon(K^N)$ be the set of associated virtual knots for all lifts by Π of K^N. If $|\Upsilon(K^N)| = 1$, i.e. all the associated virtual knots are equivalent, then the unique element υ is called the *invariant associated virtual knot*. In our notation, the main results of [4] can be restated compactly as follows.

Lemma 1. *Let* $(\mathfrak{k}_0^{\Sigma\times\mathbb{R}}, \Pi, K_0^N), (\mathfrak{k}_1^{\Sigma\times\mathbb{R}}, \Pi, K_1^N)$ *be virtual covers with invariant associated virtual knots* υ_0, υ_1*, respectively. If* $K_0^N \leftrightharpoons K_1^N$*, then* $\upsilon_0 \leftrightharpoons \upsilon_1$.

Proof. This follows from the definitions and the proof of Lemma 3 from [4]. □

1.3. **Interlude on Fibered Links.** Let $V(X)$ denote some sufficiently small tubular neighborhood of X. For a link J in $\mathbb{S}^3$, let, $N_J = \overline{\mathbb{S}^3 \backslash V(J)}$. By a fibered link, we will mean an oriented link in $\mathbb{S}^3$ having a c.c.o Seifert surface Σ_J such that the pair $(\overline{\mathbb{S}^3 \backslash V(\Sigma_J)}, \overline{\mathbb{S}^3 \backslash V(\Sigma_J)} \cap \partial N_J)$ is diffeomorphic as a pair to $(\Sigma_J \cap N_J, \partial(\Sigma_J \cap N_J)) \times \mathbb{I}$ [14]. The surface Σ_J, called a *fiber*, must be oriented so that the induced orientation on the boundary is the orientation of J. Note we may identify N_J with a mapping torus $(\Sigma_J \cap N_J) \times \mathbb{I}/\psi$, with $\psi : \Sigma_J \cap N_J \to \Sigma_J \cap N_J$ an o.p. diffeomorphism. This defines a regular covering space $\Pi_J : (\Sigma_J \cap N_J) \times \mathbb{R} \to N_J$. [1]

The Seifert surface Σ_J is of minimal genus and incompressible [14]. It is also unique in the sense that for any two minimal genus c.c.o Seifert surfaces Σ_1, Σ_2 of J, there is an ambient isotopy of $\mathbb{S}^3$ fixing J and taking Σ_1 to Σ_2 [15, 23]. This is a special property of fibered links and is not always the case for an arbitrarily chosen link.

If two fibered links have known fibers, then a new fibered link can be constructed using a generalization of the boundary connect sum called *Murasugi's sums* [6]. The resulting link is fibered and the constructed surface is a fiber. Other techniques for finding fibers Σ_J are described in [20]. More recently, there is also *Futer's method* [5]. We will take the literature on the topic as a calculus we employ to find fibers. Lastly, it is often convenient to draw fibers in *disc-band form* [1]. We may isotope these forms with *topological script* [12].

1.4. **Examples of Virtual Covers.** Consider an $m+1$ component oriented link $L = J \sqcup K$ in $\mathbb{S}^3$, where J is a fibered m component link and K is a knot in the complement N_J. The components of L are ordered, where K is the "last" component. As above, Σ_J is an oriented c.c.o Seifert surface for J. It is two-sided in $\mathbb{S}^3$, denoted red/blue in figures. The blue side is always chosen so that resting a thumb-up right hand on it reproduces the orientation of the surface and the right-handed orientation of $\mathbb{S}^3$.

There is a simple condition that guarantees a lift by Π_J. We say that K^{N_J} is in *special Seifert form (SSF) relative to* Σ_J if K^{N_J} can be decomposed into a finite number of disjoint arcs in Σ_J and a finite number of "crossings" in small disjoint embedded 3-balls $B_i \subset \text{int}(N_J)$: each $B_i \cap \Sigma_J$ is an embedded disc dividing B_i into halves, so that $K \cap B_i$ consists of two disjoint arcs in different halves of ∂B_i configured as in the model on the left side of Figure 1 (see also [4]). As Π_J is a local diffeomorphism and $\Pi_J^{-1}(\Sigma_J) \approx \Sigma_J \times \mathbb{Z}$, any K^{N_J} in SSF relative to Σ_J has a lift by Π_J. Hence we have virtual covers of the form $(\mathfrak{k}^{\Sigma_J \times \mathbb{R}}, \Pi_J, K^{N_J})$.

If K^{N_J} is in SSF relative to Σ_J, it has an invariant associated virtual knot. Note that Σ_J is a framed neat submanifold of N_J [16]. Orient the normal bundle by choosing a normal vector $\vec{v}$ at each point of Σ_J so that the surface orientation together with $\vec{v}$ gives the right-handed orientation of $\mathbb{S}^3$. Assume the SSF lies in this oriented tubular neighborhood $\Sigma_J \times (-1, 1)$ where $\Sigma_J \equiv \Sigma_J \times \{0\}$. An arc in a ball of the the SSF is *overcrossing* (resp. *undercrossing*) if it lies in $\Sigma_J \times [0, 1)$ (resp. $\Sigma_J \times (-1, 0]$). The SSF thus defines a knot diagram $[K; \Sigma_J]$ on Σ_J. By [4], the invariant associated virtual is $\kappa([K; \Sigma_J])$. See Figure 1.

For an example of virtual covers of knots K^N where N is closed, take $L = J \sqcup K$ where J

[1] Henceforth, we will frequently abuse notation and fail to distinguish between Σ_J and $\Sigma_J \cap N_J$.



FIGURE 1. (Left) A "crossing" in special Seifert form. (Right) A red knot in SSF wrt a left handed trefoil fiber and the invariant associated virtual knot.

is a fibered knot in $\mathbb{S}^3$ and K is in SSF relative to Σ_J. Let N be the manifold obtained by performing the 0-surgery on J. Then N is a fibered 3-manifold. A fiber can be obtained by gluing a disc to $\partial\Sigma_J \approx \mathbb{S}^1$. Hence K^N has an invariant associated virtual knot as above.

1.5. **The Main Theorem.** We show that the associated virtual knot can be used essentially as a link invariant. It is not an invariant in the traditional sense, but it is an invariant in the sense that if two equivalent links of the form $L = J \sqcup K$ as above have associated virtual knots, then the associated virtual knots must be equivalent. First, a lemma is needed.

Lemma 2. *If K is in SSF with respect to Σ_J and $H : \mathbb{S}^3 \to \mathbb{S}^3$ is an o.p. diffeomorphism, then $H(K)$ is in SSF with respect to $H(\Sigma_J)$ and $\kappa([K;\Sigma_J]) \leftrightharpoons \kappa([H(K);H(\Sigma_J)])$.*

Proof. An SSF is defined by embedded 3-balls B_i, discs on Σ_J, arcs on Σ_J, and "crossings" in balls B_i. These are mapped by H to 3-balls, disc/arcs on $H(\Sigma_J)$ and "crossings" in balls $H(B_i)$. $H(\Sigma_J)$ inherits its orientation from Σ_J and is a canonically oriented framed neat submanifold of $\overline{\mathbb{S}^3 \backslash V(\partial H(\Sigma_J))}$. H is o.p., so the overcrossing arcs of the SSF for K are mapped to the canonically chosen overcrossing arcs for $H(K)$ in SSF with respect to $H(\Sigma_J)$. Observe that the Gauss diagrams of $[K;\Sigma_J]$ and $[H(K);H(\Sigma_J)]$ are equivalent. □

Theorem 3 (Main Theorem)**.** *For $i = 1, 2$, let $L_i = J_i \sqcup K_i$ be oriented links with J_i a fibered link, K_i in SSF with respect to a fiber Σ_i of J_i, and υ_i the invariant associated virtual knot. If $L_1 \leftrightharpoons L_2$, then $\upsilon_1 \leftrightharpoons \upsilon_2$.*

Proof. By hypothesis, there is an o.p. diffeomorphism $H : \mathbb{S}^3 \to \mathbb{S}^3$ taking J_1 to J_2 and K_1 to K_2. Let $\Sigma_2' = H^{-1}(\Sigma_2)$. This is a minimal genus c.c.o Seifert surface for J_1. Since J_1 is a fibered link, its minimal genus Seifert surfaces are unique. Hence, there is an ambient isotopy G of $\mathbb{S}^3$ taking Σ_2' to Σ_1 and fixing J_1 pointwise. In particular, the inherited orientation on Σ_2' maps to the orientation of Σ_1. By Lemma 2, $K_1 = H^{-1}(K_2)$ is in SSF with respect to $\Sigma_2' = H^{-1}(\Sigma_2)$ and $\kappa([K_1;\Sigma_2']) \leftrightharpoons \upsilon_2$. Note also that G takes K_1 to a knot K_1' in SSF with respect to Σ_1. Again by Lemma 2, it follows that $\upsilon_2 \leftrightharpoons \kappa([K_1';\Sigma_1])$.

The ambient isotopy G is also an isotopy of the embeddings of circles K_1 and K_1'. Since K_1 and K_1' are in SSF with respect to Σ_2' and Σ_1, respectively, each level of the isotopy must lie in the interior of N_{J_1}. Hence, by the isotopy extension theorem [16], there is an ambient isotopy of N_{J_1} taking $(K_1)^{N_{J_1}}$ to $(K_1')^{N_{J_1}}$. Thus there are virtual covers $(\mathfrak{k}_1^{\Sigma_1\times\mathbb{R}}, \Pi, K_1^{N_{J_1}})$ and $((\mathfrak{k}_1')^{\Sigma_1\times\mathbb{R}}, \Pi, (K_1')^{N_{J_1}})$ and $(K_1)^{N_{J_1}} \leftrightharpoons (K_1')^{N_{J_1}}$. By Lemma 1, it follows that $\upsilon_1 \leftrightharpoons \kappa([K_1;\Sigma_1]) \leftrightharpoons \kappa([K_1';\Sigma_1]) \leftrightharpoons \upsilon_2$. □

An immediate corollary is the following strengthening of Theorem 2 from [4]. It will be used to show that some links are non-split.

Corollary 4. *Let $L = J \sqcup K$ with J a fibered link, K in SSF with respect to a fiber Σ_J of J, and υ the invariant associated virtual knot. If there is an embedded 3-ball B in $\mathbb{S}^3$ such that $K \subseteq int(B)$ and $J \subseteq \mathbb{S}^3 \backslash B$ (so that L is split), then $\upsilon \leftrightharpoons K$.*

Proof. There is an ambient isotopy moving K far away from Σ_J. Then K can be drawn in SSF with respect to some embedded 2-disc D disjoint from Σ_J. This disc can be made arbitrarily small and moved back to a disc D' on Σ_J. K' is in SSF with respect to Σ_J and clearly its invariant associated virtual $\upsilon' \leftrightharpoons K$. By Theorem 3, $\upsilon \leftrightharpoons K$. □

Example: The three component link L on the right in Figure 2 is not split. Suppose it is split, so that $\mathbb{S}^3$ decomposes into 3-balls $B_1 \cup B_2$ each containing a component of L. The black components would need to be in the same ball, say B_1, since they have linking number one. The red component K cannot be contained in B_2: K is in SSF with respect to a fiber of the Hopf link with trivial invariant associated virtual knot and $K^{\mathbb{S}^3} \leftrightharpoons 4_1$ isn't trivial. □



FIGURE 2. (Left) A non-split link $J \sqcup K$. (Middle) K in SSF with respect to a Hopf link fiber. (Right) The invariant associated virtual knot is trivial.

1.6. **Non-Invertible Links.** This section applies Theorem 3 to the detection of non-invertible links in $\mathbb{S}^3$. Denote as usual the change of orientation of all components of L by $-L$. An oriented link L is said to be *invertible* if $L \leftrightharpoons -L$ (Note: the given ordering of the components is not changed).

If $\mathfrak{k}$ is a knot diagram on a c.c.o surface, let $D_{\mathfrak{k}}$ denote any Gauss diagram of $\mathfrak{k}$. For any Gauss diagram D, let $\mathfrak{f}(D)$ denote the diagram obtained from D by changing the direction of each arrow (but not the sign).

Theorem 5. *Let $L = J \sqcup K$ be an $m+1$ component oriented link, with J an m component fibered link, Σ_J a fiber, and K a knot in SSF with respect to Σ_J. If $L = J \sqcup K$ is invertible, then $\kappa(D_{[K;\Sigma_J]}) \leftrightharpoons -\kappa(\mathfrak{f}(D_{[K;\Sigma_J]}))$.*

Proof. Recall that the orientation of Σ_J induces the given orientation on $\partial\Sigma_J = J$. A fiber Σ'_J for $-J$ may be obtained by changing the orientation of Σ_J. This also changes the framing of Σ_J; just switch the blue and red sides. Clearly, $-K$ is in SSF with respect to Σ'_J. A Gauss diagram for $[-K;\Sigma'_J]$ can be obtained from $D_{[K;\Sigma_J]}$ by changing the orientation of $D_{[K;\Sigma_J]}$ and switching each overcrossing arc in the SSF to an undercrossing arc (and vice versa). This does not change the local orientation of the crossing. Now apply Theorem 3. □

Example: The link $L = J \sqcup K$ on the left in Figure 3 is a non-split non-invertible link with invertible components, where $J \equiv$ black, $K \equiv$ red. An evident Seifert surface Σ_J in disc-band form is depicted. J is fibered and invertible: it is a connect sum of 3_1 (leftmost two bands), 5_1 (next four bands) and 3_1(rightmost two bands). As a Murasugi sum of fibers of fibered knots, Σ_J is a fiber. K is invertible as it a left handed trefoil (after some effort). A Gauss diagram D of the invariant associated virtual knot υ is on the right of the figure. The (reduced) Sawollek polynomial [19] shows that $D \not\leftrightharpoons -\mathfrak{f}(D)$:

$$\begin{aligned}\tilde{Z}_D(x,y) &= -1 + x^2 - x^3 + x^4 + \frac{x^3}{y^3} + \frac{x^4}{y^3} - y + xy + xy^2 - x^2y^2,\\ \tilde{Z}_{-\mathfrak{f}(D)}(x,y) &= -1 + x^2 - x^3 + x^4 + \frac{x^3}{y^2} - \frac{x^4}{y^2} - \frac{x}{y} + \frac{x^2}{y} - y^3 + xy^3.\end{aligned}$$

Hence, L is not invertible. The Sawollek polynomial of a classical knot is 0, so υ is non-classical and hence not equivalent to K. By Corollary 4, L is non-split. □



FIGURE 3. (Left) A non-invertible non-split link with invertible components. (Right) The invariant associated virtual knot.

The previous example can be generalized to any number of components. Whitten proved [24] that for every $m \geq 2$, there is an m component non-invertible non-split link all of whose proper sublinks are invertible. This is the inspiration behind the following result.

Corollary 6. *For every $m \geq 2$, there are infinitely many m component non-invertible non-split links $L_{m,i}$ all of whose components are invertible. Moreover, $L_{m,i} = J_{m,i,1} \sqcup \cdots \sqcup J_{m,i,m-1} \sqcup K_{m,i}$ may be chosen so that each proper sublink not containing $J_{m,i,1}$ is invertible.*

Proof. Let $\Sigma_{2,1}$ be the surface indicated in Figure 3. Construct a surface $\Sigma_{m,i}$ by taking Murasugi sums [6] with $i-1$ trefoil fibers Σ_T and $m-2$ Hopf link fibers Σ_H. The desired operations are topologically boundary connect sums of surfaces, as indicated in Figure 4. $\Sigma_{m,i}$ is an oriented fiber of the oriented fibered link $\partial\Sigma_{m,i}$ [20]. The link $\partial\Sigma_{m,i}$ has $m-2$ unknotted components and one component that is a connect sum of 5_1 and $i+1$ right handed trefoils. Set this last component to be $J_{m,i,1}$ and label the other components arbitrarily. Define $K_{m,i}$ to be the red knot K from Figure 3.

By construction, $K_{m,i}$ is in SSF with respect to $\Sigma_{m,i}$. Its invariant associated virtual knot is depicted on the right in Figure 3. Thus $L_{m,i}$ is non-invertible (see previous example). All of the components are invertible. Any proper sublink not containing $J_{m,i,1}$ is a split link that splits into a trefoil and an unlink. Such a link is invertible.

Lastly we must show that $L_{j,i}$ is non-split. If it is split, then $\mathbb{S}^3$ can be decomposed into two 3-balls B_1 and B_2 each containing at least one component of $L_{j,i}$. Linking number considerations imply that all of the components $J_{j,i,k}$ must be in the same 3-ball, say B_1. But $K_{j,i}$ cannot be in B_2 since the associated virtual knot is not classical (by Corollary 4). □



FIGURE 4. Murasugi sums used in the proof of Corollary 6.

2. Reverse Virtual Covers and the Vassiliev Filtration

2.1. **Reverse Virtual Covers.** Typically one thinks of virtual covers as associating virtual knots to *some* links. It is also fruitful to think in terms of *reverse virtual covers*: associating a link in $\mathbb{S}^3$ to a knot in a thickened surface. Let $\mathcal{V}$ denote the set of equivalence classes of oriented virtual knots. Let Γ be an oriented surface having genus g and $b \geq 1$ boundary components. Let $\mathcal{K}^{\Gamma\times\mathbb{I}}$ denote the equivalence classes of oriented knots in $\Gamma \times \mathbb{I}$. Let $\mathcal{S}$ be the set of links in $\mathbb{S}^3$ satisfying the hypotheses of Theorem 3, considered equivalent up to ambient isotopy. Then there is a well-defined map $\mathcal{I} : \mathcal{S} \to \mathcal{V}$ given by $\mathcal{I}(J \sqcup K) = v$, where v is the invariant associated virtual knot. We will show there is a map $\alpha_J : \mathcal{K}^{\Gamma\times\mathbb{I}} \to \mathcal{S}$ satisfying $\kappa = \mathcal{I} \circ \alpha_J$, where $\kappa : \mathcal{K}^{\Gamma\times\mathbb{I}} \to \mathcal{V}$ is the stabilizing map to virtual knots.

There is a fibered link J having genus g and b components (as in Figure 4). Identify N_J with the mapping torus $\Gamma \times \mathbb{I}/\psi$, where $\psi : \Gamma \to \Gamma$ is an o.p. diffeomporphism. Let Σ_J be the Seifert surface of J identified with $\Gamma \times \{1/2\}$. Assume without loss of generality that Γ is oriented so that the orientation of $\Gamma \times \mathbb{I}$ agrees with the canonical framing of Σ_J. Any knot $K^{\Gamma\times\mathbb{I}}$ may be isotoped into SSF with respect to the surface $\Gamma \times \{1/2\}$. The inclusion into the mapping torus gives a knot K' in SSF with respect to Σ_J. Define $\alpha_J(K^{\Gamma\times\mathbb{I}}) = J \sqcup K'$.

If $K_1^{\Gamma\times\mathbb{I}} \rightleftharpoons K_2^{\Gamma\times\mathbb{I}}$, then there is an ambient isotopy connecting them which acts at all times as the identity on $\partial(\Gamma \times \mathbb{I})$. Thus the isotopy extends to $\Gamma \times \mathbb{I}/\psi$. Hence $\alpha_J(K_1^{\Gamma\times\mathbb{I}}) \rightleftharpoons \alpha_J(K_2^{\Gamma\times\mathbb{I}})$ as links in $\mathbb{S}^3$ and $\alpha_J : \mathcal{K}^{\Gamma\times\mathbb{I}} \to \mathcal{S}$ is well-defined. By construction, the invariant associated virtual knot v for $J \sqcup K'$ must be $\kappa([K; \Gamma \times \{1/2\}])$. It follows that $\kappa = \mathcal{I} \circ \alpha_J$.

2.2. **The Vassiliev Filtration.** Reverse virtual covers have interesting consequences for finite-type invariants. Recall that a singular link $L_\bullet$ in a manifold N is a proper immersion of circles into N having only transverse double points in its singularity set [9]. Such links are considered equivalent up to ambient isotopy in N acting at all times as the identity on ∂N. Let $\mathcal{L}_r^N$ denote the equivalence classes of singular links with exactly r double points. For singular knots, the parallel notation $K_\bullet$, $\mathcal{K}_r^N$ will be used.



FIGURE 5. Resolution of a singular crossing.

There is a map $e_r^N : \mathcal{L}_r^N \to \mathbb{Z}[\mathcal{L}_0^N]$ which is defined as a signed sum of all possible ways to resolve the singular crossings. For $L_\bullet \in \mathcal{L}_r^N$ and $\varepsilon = (\varepsilon_1, \ldots, \varepsilon_r) \in \{1, -1\}^r$, let $L_\varepsilon \in \mathcal{L}_0^N$ denote the link obtained by resolving for all i the i-th double point according to the value of ε_i as in Figure 5. Define:

$$e_r^N(L_\bullet) = \sum_{\varepsilon \in \{\pm 1\}^r} \left(\prod_{i=1}^r \varepsilon_i \right) L_\varepsilon.$$

Set $\mathcal{J}_r^N = \left\langle e_r^N(\mathcal{L}_r^N) \right\rangle$, the subgroup of $\mathbb{Z}[\mathcal{L}_0^N]$ generated by the image of e_r^N. The *Vassiliev filtration* is the sequence of inclusions:

$$\mathbb{Z}[\mathcal{L}_0^N] \supseteq \mathcal{J}_1^N \supseteq \mathcal{J}_2^N \supseteq \cdots \supseteq \bigcap_{i=1}^\infty \mathcal{J}_r^N =: \mathcal{J}_\infty^N.$$

Two links L_1^N, L_2^N are said to be V_k^N-equivalent if $[L_1^N] - [L_2^N] \in \mathcal{J}_{k+1}^N$. By V_∞^N-equivalence, it is meant that the difference is in $\mathcal{J}_k^N$ for all k.

Theorem 7. *The map $\alpha_J : \mathcal{K}^{\Gamma \times \mathbb{I}} \to \mathcal{S}$ extends to a map $\alpha_J : \mathbb{Z}[\mathcal{K}_0^{\Gamma \times \mathbb{I}}] \to \mathbb{Z}[\mathcal{L}_0^{\mathbb{S}^3}]$ that preserves the Vassiliev filtration. In particular, $\alpha_J(\mathcal{J}_r^{\Gamma \times \mathbb{I}}) \subseteq \mathcal{J}_r^{\mathbb{S}^3}$ for all r (note here that $\mathcal{J}_r^{\Gamma \times \mathbb{I}} = \left\langle e_r^{\Gamma \times \mathbb{I}}(\mathcal{K}_r^{\Gamma \times \mathbb{I}}) \right\rangle$).*

Proof. First extend α_J by linearity to the free abelian group. Let $K_\bullet := K_\bullet^{\Gamma \times \mathbb{I}} \in \mathcal{K}_r^{\Gamma \times \mathbb{I}}$. Then

$$\alpha_J(e_r^{\Gamma \times \mathbb{I}}(K_\bullet)) = \alpha_J \left(\sum_{\varepsilon \in \{\pm 1\}^r} \left(\prod_{i=1}^r \varepsilon_i \right) K_\varepsilon \right) = \sum_{\varepsilon \in \{\pm 1\}^r} \left(\prod_{i=1}^r \varepsilon_i \right) J \sqcup K'_\varepsilon = e_r^{\mathbb{S}^3}(J \sqcup K'_\bullet),$$

since $\mathrm{int}(\Gamma \times \mathbb{I})$ embeds into $\Gamma \times \mathbb{I}/\psi$. Thus $\alpha(\mathcal{J}_r^{\Gamma \times \mathbb{I}}) \subseteq \mathcal{J}_r^{\mathbb{S}^3}$. □

Let A be an abelian group and $\nu : \mathcal{L}_0^N \to A$ a link invariant. There is an extension of ν to $\mathbb{Z}[\mathcal{L}_0^N]$ by linearity. Then ν is said to be an *A-valued finite-type invariant* of degree $\leq n$ if ν vanishes on $\mathcal{J}_{n+1}^N$. If L_1^N and L_2^N are V_n^N-equivalent, then for all abelian groups A, the A-valued finite-type invariants of degree $\leq n$ agree on L_1^N and L_2^N.

The *Vassiliev Conjecture for N* states that if all finite-type invariants of L_1^N and L_2^N agree, then $L_1^N \leftrightharpoons L_2^N$. Equivalently, the conjecture is that V_∞^N-equivalent knots are equivalent [9]. Reverse virtual covers can be used to show that if the Vassiliev conjecture is true for links in $\mathbb{S}^3$, then it is true for knots in thickened c.c.o surfaces having at least one boundary component, up to equivalence of virtual knots. More precisely, we have:

Theorem 8. *Suppose that the Vassiliev Conjecture for links in $\mathbb{S}^3$ holds. Let Γ be a c.c.o surface having at least one boundary component. If $K_1^{\Gamma\times\mathbb{I}}$ and $K_2^{\Gamma\times\mathbb{I}}$ are $V_\infty^{\Gamma\times\mathbb{I}}$-equivalent, then $\kappa(K_1^{\Gamma\times\mathbb{I}}) \leftrightharpoons \kappa(K_2^{\Gamma\times\mathbb{I}})$.*

Proof. Suppose that $K_1^{\Gamma\times\mathbb{I}}$ and $K_2^{\Gamma\times\mathbb{I}}$ are $V_\infty^{\Gamma\times\mathbb{I}}$-equivalent. Then for all k, $[K_1] - [K_2] \in \mathcal{J}_{k+1}^{\Gamma\times\mathbb{I}}$. Let J be a fibered link so that $\alpha_J : \mathcal{K}^{\Gamma\times\mathbb{I}} \to \mathcal{S}$ is well-defined. Hence for all k, $[\alpha_J(K_1)] - [\alpha_J(K_2)] \in \mathcal{J}_{k+1}^{\mathbb{S}^3}$. Since we are assuming the Vassiliev conjecture for links in $\mathbb{S}^3$, this implies that $\alpha_J(K_1) = J \sqcup K_1' \leftrightharpoons J \sqcup K_2' = \alpha_J(K_2)$. Thus, $\kappa(K_1^{\Gamma\times\mathbb{I}}) = \mathcal{I} \circ \alpha_J(K_1) \leftrightharpoons \mathcal{I} \circ \alpha_J(K_2) = \kappa(K_1^{\Gamma\times\mathbb{I}})$. Note that Theorem 3 was utilized in the last statement. □

A similar result holds for the Habiro Conjecture ([9], Conjecture 3.9). Two links in a 3-manifold M are C_k^M-equivalent if they are obtainable from one another by a finite sequence of surgeries on strict tree claspers of degree k. If two links are C_k^M-equivalent for all k, then they are said to be C_∞^M-equivalent. The Habiro Conjecture is that two links in M are equivalent if and only if they are C_∞^M-equivalent. An analogous argument to the above can be used to show that if the Habiro Conjecture holds for links in $\mathbb{S}^3$, then it holds for knots in $\Gamma \times \mathbb{I}$ up to equivalence of virtual knots.

3. Virtual Covers and Virtually Fibered Links

Previously links in $\mathbb{S}^3$ of the form $J \sqcup K$ where J is fibered have been considered. Virtual covers can also be applied to links of the form $J \sqcup K$ where J is not fibered. We will consider links where N_J has a finite index cover by $N_{\tilde{J}}$, where $\tilde{J}$ is a fibered link. Such a link is an example of a *virtually*[2] *fibered link*[22]. More generally, a virtually fibered 3-manifold is one which has a finite index cover by a fibered 3-manifold [21].



Figure 6. The links $\tilde{J}$ (Left) and J (Right) (drawn in SnapPy).

Example: We review Gabai's [7] example of a non-fibered, virtually fibered hyperbolic link. Let J be the two component link on the right hand side of Figure 6 and $\tilde{J}$ the three component

[2]The double use of the word "virtual" is admittedly confusing, but unavoidable.

link on the left hand side of Figure 6. An index two covering projection $N_{\tilde{J}} \to N_J$ can be described as follows. A "dented cube" C with a tangle removed is depicted at the bottom of Figure 7. This is a fundamental region of an index two covering projection $C \sqcup C \to C$ (see Figure 7). N_J is constructed by identifying the golden annulus and red annulus in Figure 8 and then identifying the blue and green twice punctured discs in Figure 8. The corresponding identifications on $C \sqcup C$ produce $N_{\tilde{J}}$.



FIGURE 7. Constructing Gabai's double cover $N_{\tilde{J}}$ of N_J.

The calculus described in Section 1.3 can be used to show that the surface $\Sigma_{\tilde{J}}$ in Figure 9, after an ambient isotopy of $\tilde{J}$, is a fiber for $\tilde{J}$. By [7], J is non-fibered and hyperbolic. $\square$

Let $L = J \sqcup K$ and suppose that J is a virtually fibered link admitting a finite-index covering $p : N_{\tilde{J}} \to N_J$, where $\tilde{J}$ is a fibered link. Denote the index by n. Assume also that p is regular and orientation preserving. Suppose that there is a knot $\tilde{K}$ in $N_{\tilde{J}}$ such that $p(\tilde{K}) = K$ (as functions defined on $\mathbb{S}^1$). By regularity, there are n such knots $\tilde{K} = \tilde{K}(1), \ldots, \tilde{K}(n)$. Suppose that each $\tilde{K}(i)$ is in SSF with respect to some fiber $\Sigma(i)$ of $\tilde{J}$. The *associated virtual knot spectrum* of K in N_J is the formal sum:

$$\sigma_K = \sum_{i=1}^{n} \kappa([\tilde{K}(i); \Sigma(i)]).$$

We consider σ_K as an element of $\mathbb{Z}[\mathcal{V}]$, the free abelian group generated by the equivalence classes of oriented virtual knots.

Theorem 9. *Let $K_1^{N_J}$, $K_2^{N_J}$ have associated virtual knot spectra σ_1, σ_2, respectively. If $K_1^{N_J} \leftrightharpoons K_2^{N_J}$, then $\sigma_1 = \sigma_2$.*



FIGURE 8. The identifications of the bottom cube in Figure 7 to obtain N_J.



FIGURE 9. Gabai's fiber for $N_{\tilde{J}}$.

Proof. View the ambient isotopy as an isotopy of the functions $K_i : \mathbb{S}^1 \to N_J$. The isotopy lifts to $N_{\tilde{J}}$ and connects some lift of K_1 to some lift of K_2. Again applying the isotopy extension theorem we see that these are equivalences of knots in $N_{\tilde{J}}$. Say $\tilde{K}_1(i) \leftrightharpoons \tilde{K}_2(j_i)$. Apply this to all lifts of K_1 to get a permutation of $1, \ldots, n$ defined by $i \to j_i$.

Let $\tilde{L}_k(i) = \tilde{J} \sqcup \tilde{K}_k(i)$. By the above, we have an equivalence $\tilde{L}_1(i) \leftrightharpoons \tilde{L}_2(j_i)$. By hypothesis, each $\tilde{K}_k(i)$ is in SSF with respect to some fiber $\Sigma_k(i)$. By Theorem 3, it follows that $\kappa([\tilde{K}_1(i); \Sigma_1(i)]) \leftrightharpoons \kappa([\tilde{K}_2(j_i); \Sigma_2(j_i)])$. Hence $\sigma_1 = \sigma_2$. □

Example: Let J, $\tilde{J}$ be as in Gabai's example. The covering $p : N_{\tilde{J}} \to N_J$ is regular because it has index 2. Clearly p preserves orientation. Let K be the red knot in the middle panel of Figure 10 oriented arbitrarily. We will show K is non-invertible in N_J. K has two lifts $\tilde{K}(1)$ and $\tilde{K}(2)$. $\tilde{K}(1)$ in SSF with respect to Gabai's fiber $\Sigma_{\tilde{J}}$ (top panel). The invariant associated virtual knot $\upsilon(1) = \kappa([\tilde{K}(1); \Sigma_J])$ is in the bottom left panel. This is equivalent to the non-invertible classical knot 8_{17}, oriented as $\vec{8}_{17}$. The knot $\tilde{K}(2)$ can also be put into

special SSF with respect to Σ_J (after some effort). The associated virtual knot $\upsilon(2)$ is also $\vec{8}_{17}$. Hence $\sigma_K = 2 \cdot \vec{8}_{17}$. But $\sigma_{-K} = 2 \cdot (-\vec{8}_{17})$. Hence $K \neq -K$ in N_J. Note however that K is a trefoil in $\mathbb{S}^3$, and hence is invertible in $\mathbb{S}^3$ (bottom right panel). □

3.1. **Acknowledgments.** This work was part of a Spring 2015 sabbatical from Monmouth University. For help and encouragement, thanks to A. Kaestner, K. Orr, and R. Todd.

References

[1] G. Burde and H. Zieschang. *Knots*, volume 5 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, second edition, 2003.

[2] J. S. Carter, S. Kamada, and M. Saito. Stable equivalence of knots on surfaces and virtual knot cobordisms. *J. Knot Theory Ramifications*, 11(3):311–322, 2002. Knots 2000 Korea, Vol. 1 (Yongpyong).

[3] M. Chrisman and A. Kaestner. Virtual covers of links II. *in preparation*.

[4] M. W. Chrisman and V. O. Manturov. Fibered knots and virtual knots. *J. Knot Theory Ramifications*, 22(12):1341003, 23, 2013.

[5] D. Futer. Fiber detection for state surfaces. *Algebr. Geom. Topol.*, 13(5):2799–2807, 2013.

[6] D. Gabai. The Murasugi sum is a natural geometric operation. In *Low-dimensional topology (San Francisco, Calif., 1981)*, volume 20 of *Contemp. Math.*, pages 131–143. Amer. Math. Soc., Providence, RI, 1983.

[7] D. Gabai. On 3-manifolds finitely covered by surface bundles. In *Low-dimensional topology and Kleinian groups (Coventry/Durham, 1984)*, volume 112 of *London Math. Soc. Lecture Note Ser.*, pages 145–155. Cambridge Univ. Press, Cambridge, 1986.

[8] M. Goussarov, M. Polyak, and O. Viro. Finite-type invariants of classical and virtual knots. *Topology*, 39(5):1045–1068, 2000.

[9] K. Habiro. Claspers and finite type invariants of links. *Geom. Topol.*, 4:1–83 (electronic), 2000.

[10] R. Hartley. Identifying noninvertible knots. *Topology*, 22(2):137–145, 1983.

[11] N. Kamada and S. Kamada. Abstract link diagrams and virtual knots. *Journal of Knot Theory and its Ramifications*, 9:93–106, 2000.

[12] L. H. Kauffman. *On knots*, volume 115 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1987.

[13] L. H. Kauffman. Virtual knot theory. *European J. Combin.*, 20(7):663–690, 1999.

[14] A. Kawauchi. *A survey of knot theory*. Birkhäuser Verlag, Basel, 1996. Translated and revised from the 1990 Japanese original by the author.

[15] T. Kobayashi. Uniqueness of minimal genus Seifert surfaces for links. *Topology Appl.*, 33(3):265–279, 1989.

[16] A. A. Kosinski. *Differential manifolds*, volume 138 of *Pure and Applied Mathematics*. Academic Press, Inc., Boston, MA, 1993.

[17] V. A. Krasnov and V. O. Manturov. Graph-valued invariants of virtual and classical links and minimality problem. *J. Knot Theory Ramifications*, 22(12):1341006, 14, 2013.

[18] G. Kuperberg. What is a virtual link? *Algebraic and Geometric Topology*, 3:587–591, 2003.

[19] J. Sawollek. On Alexander-Conway Polynomials for Virtual Knots and Links. *arXiv:9912173[math.GT]*, December 1999.

[20] J. R. Stallings. Constructions of fibred knots and links. In *Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 2*, Proc. Sympos. Pure Math., XXXII, pages 55–60. Amer. Math. Soc., Providence, R.I., 1978.

[21] W. P. Thurston. Three-dimensional manifolds, Kleinian groups and hyperbolic geometry. *Bull. Amer. Math. Soc. (N.S.)*, 6(3):357–381, 1982.

[22] G. S. Walsh. Great circle links and virtually fibered knots. *Topology*, 44(5):947–958, 2005.

[23] W. Whitten. Isotopy types of knot spanning surfaces. *Topology*, 12:373–380, 1973.

[24] W. C. Whitten, Jr. On noninvertible links with invertible proper sublinks. *Proc. Amer. Math. Soc.*, 26:341–346, 1970.



FIGURE 10. The knot K in N_J (middle, red), the knot $\tilde{K}(1)$ in $N_{\tilde{J}}$ (top, red), associated virtual knot $v(1) \rightharpoonup 8_{17}$ (bottom left), and K depicted as a left handed trefoil in $\mathbb{S}^3$ (bottom right).